



Allometric Formulas to Predict *Haematostaphis barteri* Biomass Above and Below Ground in Cameroon

Dr. Awé Djongmo Victor^{1*} and Prof. Noiha Noumi Valéry²

¹Department of Plant Production, Higher Institute of Agriculture, Wood, Water Resources and Environment of Belabo, University of Bertoua, P.O. Box 60, Belabo, Cameroon

²Department of Life Science, Higher Teacher Training College of Bertoua, University of Bertoua, Cameroon



Abstract

This study used a destructive approach to develop species-specific allometric equations for *H. barteri* in Cameroon. A sample of 30 individuals was chopped, measured, and weighed, and six prediction models were created for leaves, branches, trunks, roots, and total above-ground biomass. The best models were chosen using Akaike's information criterion, residual standard error, root mean square error, and adjusted coefficients of determination. The total biomass above ground was $\ln(B) = 2.832 + 0.981 * \ln(\text{dbh}^2 \times H \times \text{WD}) * (1.0008)$; the trunks were $\ln(B) = 1.101 + 0.106 * \ln(\text{dbh} \times \text{WD}) * (1.0003)$; the branches were $\ln(B) = 0.065 + 0.010 * \ln(\text{dbh} \times \text{WD}) * (1.0008)$; the leaves were $\ln(B) = 0.065 + 0.010 * \ln(\text{dbh}^2 \times H) * (1.0008)$, and the roots were $\ln(B) = 0.758 + 0.035 * \ln(\text{dbh}) * (1.0009)$. The equations offer important information for scaling up biomass estimations and evaluating carbon sequestration potential in *H. barteri* stands in Cameroon.

Keywords: *Haematostaphis barteri*; Cameroon, Biomass, Allometric Formulas

Introduction

Haematostaphis barteri Hook. F. is a forest tree species with regional significance found in savannas of Guinea and Sudan in tropical Africa [1-3]. Its fruits are marketed fresh in local marketplaces and consumed by people in North Cameroon, Togo, and northern Cameroon. *H. barteri* contains protein, vitamins, and trace elements in its fruits and leaves. Its natural range is limited to tropical Africa, including Ghana, Togo, Benin, Nigeria, Cameroon, and Chad [4]. Forests play a significant role in the global carbon cycle, and REDD+ programs aim to estimate carbon stored in forests [5]. Local allometric models are essential for estimating biomass in various land uses, such as nitrogen cycle, energy, and environmental impacts. Forests are considered a means of mitigating climate change [5].

Wood basic density (WD) is a crucial explanatory element for biomass estimation, calculated from wood samples by dividing dry mass by green volume [5-12]. Tree dbh and ht are often used as explanatory variables in biomass allometric models [5, 7, 13]. However, the dbh range, unequal distribution, and ecological zones limit their applications in Africa [14]. Some sub-Saharan African nations rely on pantropic models to estimate local AGB, but these models cannot solve the increasing demand for local allometric models [15, 16].

Despite its importance, little information exists about its potential for carbon storage. Understanding biomass could help rural populations become more resilient to climate change and aid carbon sequestration initiatives like the REDD + Mechanism biomass [17-22]. The study aims to create allometric formulas to calculate the aboveground and root biomass of *H. barteri* in Cameroon's Sudano-Sahelian zone.

Materials and Methods

Research site

The research was conducted in Central Africa, namely in the North region of Cameroon. This area lies between latitudes 9° 18'N and 8° 10'N and longitudes 13° 23'E and 12° 16'E [23] (Figure 1). Between the Adamawa Plateau to the south and the Mandara Mountains (1442 m) to the north, the relief is a sizable pedi-plain. There are two distinct seasons in the Sudano-Sahelian climate: a six-month dry season (November–May) and a six-month rainy season (June–October) [24]. Between August and March, the average monthly temperature ranged from 26°C to 40°C. The ferruginous

OPEN ACCESS

*Correspondence:

Dr. Awé Djongmo Victor, Department of Plant Production, Higher Institute of Agriculture, Wood, Water Resources and Environment of Belabo, University of Bertoua, P.O. Box 60, Belabo, Cameroon,
E-mail: awevictor920@gmail.com

Received Date: 08 Sep 2025

Accepted Date: 16 Sep 2025

Published Date: 18 Sep 2025

Citation:

Victor AD, Valéry NN. Allometric Formulas to Predict *Haematostaphis barteri* Biomass Above and Below Ground in Cameroon. WebLog J Environ Sci. wjets.2025.i1801. <https://doi.org/10.5281/zenodo.17282572>

Copyright© 2025 Dr. Awé Djongmo Victor. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

kind of soil is distinguished by its poor cation exchange capacity and acidity (pH = 5.5–6) [25]. Around the villages, the vegetation is a clear and degraded Sudanian savannah, with shrubby vegetation [26].

Data Collection

The study used a direct method to investigate *H. barteri* trees, selecting individuals based on availability and absence of signs of human abuse. Circumferences were measured and three diameter classes 5–10 cm, 10–20 cm, and 20–30 cm were chosen, with thirty trees in the North Cameroon region marked.

Ten individuals were divided into classes and separated into three sections: leaves, branches, and trunks. Roots were excavated to determine biomass. Large branches, trunks, and roots were chopped into small pieces, bagged, and weighed. Samples were dried in an oven set to 70°C for leaves and 105°C for trunk, branches, and root discs. The following formula will be used to determine the water content of the leaf, branch, and trunk samples:

$$WC (\%) = ((WM - DM) / DM) * 100 \text{ [27, 28]}$$

where WM and DM stand for the sample's wet mass (Kg) and dry mass (Kg), respectively, and WC is the water content of the samples expressed as a percentage. The following formula was used to determine the total dry masses of the fractions based on the samples' water content: $100 * TWM / (100 + WC) = TDM$ [27, 28], TDM stands for total dry mass. The total wet mass is TWM (Kg). Known as biomass, the total dry masses are measured in kilograms (Kg).

Wood samples from trunks and branches were harvested to calculate *H. barteri*'s wood density. They were weighed, labeled, and stored in a laboratory. To calculate volume, samples were soaked in water and calibrated on a balance following Archimedes' principle [29]. Samples were dried in an oven at 105°C for 72 hours, then weighed every six hours until a consistent weight was achieved, with data stored after achieving complete evaporation of water [22, 30]. Using the dry weight to volume ratio and the calculation from [29] below, the density of the wood was determined: $WD_i = M_i / V_i$, where WD_i is the wood density i , M_i is the species i 's dry weight, and V_i is its water content. We determined the root:shoot ratio (RS) for each tree by simply dividing the values of leaves, branches, trunk, and AGB by the matching BGB value after estimating AGB and BGB as previously said.

Data Analysis

The leaf, trunk, branch, root biomass, and total AGB of *H. barteri* were found to be allometrically related to the tree's physical characteristics, including height (H), wood density (WD), and diameter at breast height (dbh) [27]. The power model and the polynomial model are the two types of models commonly used in the literature to forecast these biomasses [7]. Since the polynomial model frequently displays aberrant behavior outside of its region of validity, the power model was employed in this investigation [27].

$B = a * Db$ is the mathematical formula frequently used to modify biomass [27, 31], where B is the biomass, D is the diameter, and a , b are the regression coefficients. The formula is frequently altered using the logarithmic transformation using the relation $\ln(B) = a + b * \ln(D)$ [21, 28, 31] in order to account for the heteroskedasticity of the data [21, 27]. The following six models were used to fit the biomasses in this investigation [5, 7, 31]:

$$\ln(B) = a + b * \ln(dbh) \quad (1)$$

$$\ln(B) = a + b * \ln(dbh * \rho) \quad (2)$$

$$\ln(B) = a + b * \ln(dbh^2 * H) \quad (3)$$

$$\ln(B) = a + b * \ln(dbh^2 * H * \rho) \quad (4)$$

$$\ln(B) = a + b * \ln(dbh) + c * \ln(H) \quad (5)$$

$$\ln(B) = a + b * \ln(dbh) + c * \ln(H) + d * \ln(\rho) \quad (6)$$

In this case, B stands for biomass (kg), Dbh for tree diameter, H for total height (m), ρ or WD for wood density, and a , b , c , and d for regression coefficients.

A correction is consequently required, which entails multiplying the estimated Biomass by a correction factor (CF), which is computed as follows: The logarithmic processing of the data typically results in a bias in the calculation of Biomass [31]. [21, 27] $CF = \exp(RSE^2/2)$; the CF is always bigger than 1 [31]. The models' accuracy and robustness in estimating above-ground biomass were evaluated using four criteria [31]. They are listed in priority order: i) Adjusted R^2 , or adjusted coefficient of determination, where SRS stands for sum of residual squares and STS for sum of total squares. ii) The Akaike Information Criterion, or AIC, was calculated using the formula below: $AIC = -2\ln(L) + 2p$, where p is the total number of model parameters and L is the "Likelihood," or probability, at which the predicted model is accurate to the unknown true. iii) Residual standard error, or RSE, is calculated as follows: $RSE = \ln(AGB \text{ obs}) - \ln(AGB \text{ pred})$, where $AGB \text{ obs}$ is the measured above-ground biomass, $Pred \text{ AGB}$ is the predicted above-ground biomass. iv) Root mean squared error, or RMSE:

$$RMSE = \sqrt{\frac{\sum (\ln(AGB_{obs}) - \ln(AGB_{pred}))^2}{n - k}}$$

in which: n : the total number of observations that the model uses;

$AGB \text{ obs}$: Above-ground biomass measurement;

Predicted above-ground biomass, or $Pred \text{ AGB}$.

K : the total number of model parameters.

A number of statistical characteristics were taken into consideration while choosing the optimum allometrics created for predicting the biomass of the various *H. barteri* components. Therefore, the better the model, the lower the mean square error (RMSE), residual standard error (RSE), Akaike information criterion (AIC), and the strong adjusted R^2 [31]. R i386 3.1.2 and Excel 2020 were used to perform the statistical analyses.

Results

Height, Diameter, Wood Density, and Leaves, Branches, Trunk, Roots, Total Aboveground Biomass, and Measurable Destructive Biomass Parameters Pearson's Correlation

Table 1 shows the distribution of biomasses and dendrometric characteristics in a forest. The dbh ranged from 5.09 to 29.93 cm, with heights from 3.5 to 9.5 meters. The wood density ranged from 0.390 to 0.670 g/cm³. The biomass's AGB ranged from 164.02–298 kg, with trunks acquiring more biomass than other compartments. The biomass of leaves, trunks, branches, roots, total aboveground biomass, and H showed a positive association with the dbh , while the dbh and wood density showed a positive but non-significant link.

Regression Relationship between Diameter-Height

Individual height and diameter have been found to correlate




Figure 1: Study area's geographic location within the North Cameroon Region.




Figure 2: Diameter-height regression connection.

significantly ($r^2 = 0.785$; $P < 0.001$) (Figure 2).

Allometric Equation Development and Modeling

The regression coefficients of the examined models are statistically significant, with coefficients varying between compartments. The adjusted coefficients of determination range from 94.58 to 99.87 percent, indicating positive and significant connections between biomass and various characteristics (Table 2).

Choosing the Best Models

Leaf biomass was best predicted using $\text{dbh}^2 \times H$, with adjusted R^2 (98.98%) and RSE (0.201 kg). Branch and trunk biomass had higher adjusted R^2 values and lower RSE.

The best prediction for total above-ground biomass is achieved

using $\text{dbh}^2 \times H \times \text{WD}$, with an adjusted R^2 of 99.87%, RSE of 0.31 kg, AIC of 10.01, and RMSE of 0.143 kg. Table 3 displays these optimal equations, and their modifications are displayed in Figures 3 (a, b, c, d, e).

The proportions of total above-ground biomass, biomass from other above-ground compartments, and biomass from roots

The dug individuals' root biomass varied between 40.90 and 95.02 kg. The average root biomass to various tree parts ratios were then calculated (Table 4). Root biomass to other tree components has an average ratio between 0.29 and 1.66. Nonetheless, the overall biomass is the most significant biomass in the calculations.

Discussion

Accurate biomass models are crucial for estimating forest carbon stocks and providing verifiable information to decision-makers [27]. The diameter-height relationship indicates species' ecological growth conditions [28]. The height of a tree is predicted by its diameter, indicating variations among species due to their specific structures, competition, or environmental conditions affecting growth rate [32].

The study calculates an average base wood density of 0.53 g/cm^3 for *H. barteri*, similar to other species like *Anacardium occidentale* [33] and *Mangifera indica* [34]. However, this density is lower than previous estimates for *Tectona grandis* plantations [35]. In addition, [36] reported an average value of 0.54 g/cm^3 from 123 species in the tropical forest of Panama and [37] reported an average value of 0.60 g/cm^3 for 470 species from tropical America. The differences between the study and database values suggest potential errors in biomass estimates. Field measurements can improve allometric equation precision.

The study found that trunk biomass contributed the most to aboveground biomass, accounting for 52.35%, while leaf biomass made up the smallest part, aligning with previous literature [38, 39]. The reason for this is that the leaves are pushed onto the younger branches rather than the older ones [40]. The study involved 30 individuals, a variable sample size in allometric model development, considering resources and time allocated to the study [27]. Some allometric biomass equations have been constructed from a limited number of individuals, 26 trees [35]; 20 trees [41]; 38 trees [42]; 17 trees [27]; 20 trees [28]. Others incorporate very few large diameter trees, 1 to 79 cm in diameter [17]. Dendrometry range is a condition for model use, variable depending on objectives, species, populations,

Table 1: Height, wood density, leaves, branches, trunks, roots, total aboveground biomass, belowground biomass, and quantifiable damaging biomass factors are correlated.

Person correlation						
Item	dbh	H	WD or ρ	Mean (CV)	Range	
Leaves (kg)	0.875**	0.654**	0.265ns	40.90 (39.95)	20.93-60.88	
Branches (kg)	0.954**	0.503**	0.554**	85.52 (41.68)	64.68-106.36	
Trunk (kg)	0.812**	0.696**	0.587**	104.58 (52.35)	78.41-130.76	
TAGB (kg)	0.987**	0.785**	0.502**	2 31(133.98)	164.02-298	
Roots (kg)	0.765**	0.453ns	0.328ns	67.96 (54.12)	40.90-95.02	
dbh (cm)	1	0.785**	0.213ns	17.51 (24.84)	5.09-29.93	
H (m)	0.785**	1	0.276ns	6.50 (10.48)	3.50-9.50	
WD (g/cm^3)	0.213ns	0.276ns	1	0.530 (28)	0.390-0.670	

CV: Coefficient of variation, ** $P < 0.01$, ns: $P > 0.05$, Diameter (D), Height (H), wood density (WD), Aboveground biomass (AGB)

Table 2: Dry biomass prediction allometric equation models based on various *H. barteri* components.

Compartment	Allometric models									
	Regressions of coefficient's				Performance of model					
	a (sd)	b (sd)	c (sd)	d (sd)	Adj.R ² (%)	RSE	AIC	RMSE	P	CF
Leaf biomass										
$\ln(B) = a + b \times \ln(dbh)$	0.845(0.05)	0.203(0.03)			95.96	0.251	15.94	0.128	<0.001	1.0004
$\ln(B) = a + b \times \ln(dbh \times WD)$	1.105(0.14)	0.103(0.02)			95.88	0.228	13.27	0.124	<0.001	1.0001
$\ln(B) = a + b \times \ln(dbh^2 \times H)$	2.005(0.16)	1.035(0.01)			98.98	0.201	10.94	0.101	<0.001	1.0002
$\ln(B) = a + b \times \ln(dbh^2 \times H \times WD)$	0.205(0.02)	0.630(0.04)			95.97	0.244	16.28	0.124	<0.001	1.0006
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H)$	0.518(0.08)	0.205(0.03)	0.020(0.02)		95.98	0.254	18.94	0.141	<0.001	1.0003
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H) + d \times \ln(WD)$	0.408(0.03)	0.105(0.02)	0.010(0.01)	0.005(0.00)	94.58	0.258	14.26	0.154	<0.001	1.0005
Branch biomass										0.181
$\ln(B) = a + b \times \ln(dbh)$	0.404(0.02)	0.006(0.00)			97.98	0.204	14.14	0.120	<0.001	1.0006
$\ln(B) = a + b \times \ln(dbh \times WD)$	0.065(0.01)	0.010(0.00)			99.65	0.102	12.94	0.100	<0.001	1.0008
$\ln(B) = a + b \times \ln(dbh^2 \times H)$	0.657(0.03)	0.202(0.01)			98.08	0.118	15.28	0.114	<0.001	1.0001
$\ln(B) = a + b \times \ln(dbh^2 \times H \times WD)$	0.708(0.04)	0.307(0.02)			96.07	0.218	15.94	0.134	<0.001	1.0005
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H)$	0.902(0.05)	0.408(0.03)	0.005(0.00)		95.16	0.144	19.20	0.152	<0.001	1.0002
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H) + d \times \ln(WD)$	0.705(0.04)	0.205(0.01)	0.201(0.01)	0.102(0.00)	95.73	0.285	17.94	0.178	<0.001	1.0003
Trunk biomass										
$\ln(B) = a + b \times \ln(dbh)$	0.706(0.02)	0.326(0.04)			97.56	0.258	15.65	0.184	<0.001	1.0002
$\ln(B) = a + b \times \ln(dbh \times WD)$	1.101(0.04)	0.106(0.01)			97.88	0.212	11.01	0.112	<0.001	1.0003
$\ln(B) = a + b \times \ln(dbh^2 \times H)$	1.306(0.05)	0.406(0.02)			97.58	0.268	13.29	0.128	<0.001	1.0006
$\ln(B) = a + b \times \ln(dbh^2 \times H \times WD)$	1.017(0.01)	0.506(0.03)			97.36	0.344	16.19	0.134	<0.001	1.0005
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H)$	2.013(0.07)	0.702(0.05)	0.068(0.01)		97.52	0.318	15.05	0.120	<0.001	1.0008
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H) + d \times \ln(WD)$	3.085(0.09)	0.004(0.00)	0.002(0.00)	0.001(0.00)	97.63	0.254	13.53	0.194	<0.001	1.0004
Total aboveground biomass										
$\ln(B) = a + b \times \ln(dbh)$	1.876(0.05)	0.406(0.02)			98.98	0.312	17.59	0.189	<0.001	1.0007
$\ln(B) = a + b \times \ln(dbh \times WD)$	1.986(0.06)	0.657(0.03)			98.75	0.328	13.59	0.152	<0.001	1.0005
$\ln(B) = a + b \times \ln(dbh^2 \times H)$	1.806(0.04)	1.030(0.05)			98.74	0.344	12.01	0.148	<0.001	1.0001
$\ln(B) = a + b \times \ln(dbh^2 \times H \times WD)$	2.832(0.08)	0.981(0.04)			99.87	0.310	10.01	0.143	<0.001	1.0008
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H)$	2.302(0.07)	0.068(0.02)	0.062(0.02)		98.94	0.344	16.32	0.180	<0.001	1.0004
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H) + d \times \ln(WD)$	1.089(0.03)	0.049(0.01)	0.004(0.00)	0.001(0.00)	98.72	0.378	13.05	0.179	<0.001	1.0003
Root biomass										
$\ln(B) = a + b \times \ln(dbh)$	0.758(0.02)	0.035(0.01)			97.96	0.154	14.15	0.119	<0.001	1.0009
$\ln(B) = a + b \times \ln(dbh \times WD)$	2.206(0.08)	0.495(0.03)			95.95	0.180	18.61	0.152	<0.001	1.0001
$\ln(B) = a + b \times \ln(dbh^2 \times H)$	1.806(0.05)	0.205(0.02)			95.78	0.244	17.52	0.148	<0.001	1.0006
$\ln(B) = a + b \times \ln(dbh^2 \times H \times WD)$	0.721(0.02)	0.208(0.03)			96.78	0.218	18.15	0.184	<0.001	1.0004
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H)$	0.550(0.01)	0.062(0.01)	0.004(0.01)		96.06	0.264	15.61	0.180	<0.001	1.0008
$\ln(B) = a + b \times \ln(dbh) + c \times \ln(H) + d \times \ln(WD)$	1.008(0.04)	0.452(0.03)	0.043(0.00)	0.002(0.00)	97.43	0.268	16.52	0.179	<0.001	1.0003

The coefficients at $p < 0.001$ are significantly different from zero; Standard deviation (sd), Coefficient of regression model (a, b, c and d), Diameter at breast height (dbh), Height (H), wood density (WD), Biomass (B), logarithm (ln), adjusted coefficient of determination (adj. R²), correction factor (CF), residual standard error (RSE), Root mean squared error (RMSE) and Akaike information criteria (AIC)

and sampling plans [43]. Logarithmic transformation is used to reduce residual heterogeneity and approach linearity [44].

Allometric equations were developed using breast height, height, and wood density, considering Akaike information criterion, residual standard error, and root mean square error. Five models were selected, with Eq.4 being the best for determining total aboveground biomass and Eq. 3 for leaf biomass estimation [31]. This result corroborates those of [27] and [28] who showed respectively that the best model

for the prediction of the foliar biomass of 17 feet of *Daniella oliveri* and 20 feet of *Faidherbia albida* in Cameroon is the one taking into account the diameter squared multiplied by the height ($dbh^2 \times H$). The optimal model for estimating branch and trunk biomass is model Eq. 2, which has the lowest AIC, RMSE, and RES values, and the highest adjusted coefficient of determination for root biomass, considering only diameter. The dbh alone is a good predictor for estimating the roots biomass. This result is similar to that of [27] who worked on 17 individuals of *Daniellia oliveri* in Cameroon.

Table 3: Top allometric model selections by compartment.

Compartment biomass (kg)	Allometric models	R² (%)	RSE	RMSE	AIC
Leaves	$\ln(B) = 2.005 + 1.035 \cdot \ln(\text{dbh}^2 \times H) \cdot (1.0002)$	98.98	0.201	0.101	10.94
Branch	$\ln(B) = 0.065 + 0.010 \cdot \ln(\text{dbh} \times WD) \cdot (1.0008)$	99.65	0.102	0.100	12.94
Trunk	$\ln(B) = 1.101 + 0.106 \cdot \ln(\text{dbh} \times WD) \cdot (1.0003)$	97.88	0.212	0.112	11.01
AGB	$\ln(B) = 2.832 + 0.981 \cdot \ln(\text{dbh}^2 \times H \times WD) \cdot (1.0008)$	99.87	0.310	0.143	10.01
Roots	$\ln(B) = 0.758 + 0.035 \cdot \ln(\text{dbh}) \cdot (1.0009)$	97.96	0.154	0.119	14.15

Diameter at breast height (dbh), Height (H), wood density (WD), Biomass (B), logarithm (ln), adjusted coefficient of determination (adj. R²), residual standard error (RSE), Root mean squared error (RMSE) and Akaike information criteria (AIC)




Figure 3: Regression of the best models chosen. (a) ln (leaves biomass) and ln(dbh²×H), (b) ln (branches biomass) and ln (dbh×WD), (c) ln (trunk biomass) and ln (dbh×WD), (d) ln (roots biomass) and ln (dbh), (e) ln (AGB) and ln (dbh²×H ×WD).

Table 4: Average ratios by compartment.

Ratios	Roots biomass /Leaves biomass	Roots biomass /Branches biomass	Roots biomass/ Trunk biomass	Roots biomass/ AGB
Mean (sd)	1.66 (0.83)	0.79 (0.32)	0.64 (0.21)	0.29 (0.12)

Standard deviation (sd), Aboveground biomass (AGB)

Logarithmic CF is a statistical tool for eliminating biases, but it's generally small compared to biomass estimate variation, making it omitted [27, 32]. The study found low CF values for all biomass equations, indicating minimal errors when fitting logarithmic transformations to biomass data.

The ratio established in our study is in line with other results found in the literature such as those of [27]. We did not find results for *H. barteri* in particular, but for woody plants in general, [7] determined a ratio of 0.28 which is close to the value found in our study. The small difference observed could be due, on the one hand, to the architectural forms [27] between the species used in this study and those studied by [27] and on the other hand, to the environmental conditions which are different in the two studies.

Conclusion

This study developed allometric relationships for estimating biomass for socio-economically important species like *H. barteri* in Central Africa's Sudano-Sahelian savannah zone of Cameroon. The

study used data from thirty trees to build allometric mathematical models based on diameter, height, and wood density. The models showed varied performance, with the best model for predicting total above-ground biomass being $2.832 + 0.981 \cdot \ln(\text{dbh}^2 \times H \times WD) \cdot (1.0008)$ and the roots were $\ln(B) = 0.758 + 0.035 \cdot \ln(\text{dbh}) \cdot (1.0009)$. The models can also serve as benchmarks for formulating conservation strategies. However, new allometric models should be used cautiously when estimating biomass of trees outside their range of data and site conditions.

Acknowledgements

Authors thank all the referred whose contributions have been very significant for the improvement of this study.

References

- Biaou SSH, Moutouama JK, Dan BSC, Amahowé OI, Moutouama FT, Natta AK. Uses of *Haematostaphis barteri* Hook.f. among the Waaba and Bètammaribè in North-Benin and impact on the species vulnerability. *International Journal of Biodiversity and Conservation*. 2017; 9(5): 146-157. DOI:10.5897/IJBC2016.1063
- Arbonnier M. *Trees, Shrubs and Vines of West Africa* (4th edition). Éditions Quae, Hors Collection, 02689NUM. 2019;779.
- Yougouda H, Balna J, Souare K. Structure écologique et production fruitière de *Haematostaphis barteri* Hook. F en la zone sahélienne du Cameroun. *Journal of Applied Biosciences*. 2018; 130: 13232-13243. DOI:10.4314/jab.v130i1.10

4. Sourou BNK. Importance socio-économique et caractérisation structurale, morphologique et génétique moléculaire de *Haematostaphis barteri* Hook F. (la prune rouge) au Bénin. Mémoire de thèse de Doctorat en Sciences Agronomiques de l'Université de Parakou, Bénin. 2017; 125.
5. Segura MA, Luis MA, Hernán JA. Allometric models to estimate aboveground biomass of small trees in wet tropical forests of colombian pacific area. *Rev. Árvore*. 2018; 42(2): 1806-9088. Available: <http://dx.doi.org/10.1590/1806-90882018000200009>.
6. Chave J, Andalo C, Brown S, Cairns MA, Chambers JQ, Eamus D, Fölster H, Fromard F, Higuchi N, Kira T, Lescure JP, Nelson BW, Ogawa H, Puig H, Riéra B, Yamakura T. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*. 2005; 145: 87-99.
7. Henry M, Besnard A, Asante WA, Eshun J, Adu-Bredu SA, Valentini R, Bernoux M, Saint-André L. Wood density, phytomass variations within and among trees, and allometric equations in a tropical rainforest of Africa. *For Ecol Manag*. 2010; 260: 1375-1388.
8. Chave J, Rejou MM, Burquez A, Chidumayo E, Colgan MS, Delitti WBC, Duque A, Eid T, Fearnside PM, Goodman RC, Henry M, Martinez-Yrizar A, Mugasha WA, Muller-Landau HC, Mencuccini M, Nelson BW, Ngomanda A, Nogueira EM, Ortiz-Malavassi E, Pe'lissier R, Ploton P, Ryan CM, Saldarriaga JG, Vieilledent G. Improved allometric models to estimate the aboveground biomass of tropical trees. *Glob Change Biol*. 2014; 20:3 177-3190. <https://doi.org/10.1111/gcb.12629>
9. Taerøe A, Nord-Larsen T, Stupak I, Raulund-Rasmussen K. Allometric Biomass, Biomass Expansion Factor and Wood Density Models for the OP42 Hybrid Poplar in Southern Scandinavia. *Bioenerg. Res*. 2015; 8: 1332-1343. DOI: 10.1007/s12155-015-9592-3
10. Mugasha WA, Mwakalukwa EE, Luoga E, Malimbwi RE, Zahabu E, Silayo DS, Sola G, Crete P, Henry M, Kashindye A. Allometric Models for Estimating Tree Volume and Aboveground Biomass in Lowland Forests of Tanzania. *International Journal of Forestry Research*. 2016; (4): 1-13. DOI:10.1155/2016/8076271
11. Aabeyir R, Adu-Bredu S, Agyare WA, Weir MJC. Allometric models for estimating aboveground biomass in the tropical woodlands of Ghana, West Africa. *Forest Ecosystems*. 2020; 7: 41. Available:<https://doi.org/10.1186/s40663-020-00250-3>
12. Mukuralinda A, Kuyah S, Ruzibiza M, Ndoli A, Nabahungu NL, Muthuri C. Allometric equations, wood density and partitioning of aboveground biomass in the arboretum of Ruhunde, Rwanda. *Trees, Forests and People*. 2021; 3: 100050. Available:<https://doi.org/10.1016/j.tfp.2020.100050>
13. Ilyas S. Allometric equation and carbon sequestration of *Acacia mangium* Willd. in coal mining reclamation areas. *Civil and Environmental Research*. 2013; 3(1): 8-16.
14. Youkhana AH, Ogoshi RM, Kiniry JR, Meki MN, Nakahata MH, Crow SE. Allometric models for predicting aboveground biomass and carbon stock of tropical perennial C4 grasses in Hawaii. *Front. Plant Sci*. 2017; 8: 650. DOI: 10.3389/fpls.2017.00650
15. Ekoungoulou R, Liu X, Loumeto J, Ifo SA. Above-and belowground biomass allometrics for carbon stocks estimation in secondary forest of Congo. *J. Environ. Sci. Toxicol. Food Technol*. 2014; 8: 9-20.
16. Fard ME, and Heshmati AH. Predication of biomass of three perennial range grasses using dimensional analysis. *Middle East J. Sci. Res*. 2014; 21: 1520-1525. DOI: 10.5829/idosi.mejsr.2014.21.09.21702
17. Djomo AN, Adamou I, Joachim S, Gode G. Allometric equations for biomass estimations in Cameroon and pan moist tropical equations including biomass data from Africa. *Forest Ecology and Management*. 2010; 260: 1873-1885.
18. Djomo AN, Knohl A. & Gravenhorst G. Estimations of total ecosystem carbon pools distribution and carbon biomass current annual increment of a moist tropical forest. *For. Ecol. Manage*. 2011; 261(8): 1448-1459. <https://doi.org/10.1016/j.foreco.2011.01.031>
19. Ouédraogo K, Dimobe K, Thiombiano A. Allometric models for estimating aboveground biomass and carbon stock for *Diospyros mespiliformis* in West Africa. *Silva Fennica*. 2020; 54(1): 2242-4075. <https://doi.org/10.14214/sf.10215>
20. Picard N, Rutishauser E, Ploton P, Ngomanda A, Henry M. Should tree biomass allometry be restricted to power models? *For Ecol Manag*. 2015; 353: 156-163. <https://doi.org/10.1016/j.foreco.2015.05.035>
21. Djomo AN, Picard N, Fayolle A, Henry M, Ngomanda A, Ploton P, McLellan J, Saborowski J, Adamou I, Lejeune P. Tree allometry for estimation of carbon stocks in African tropical forests. *Forestry*. 2016; 89: 446-455. <https://doi.org/10.1093/forestry/cpw025>
22. Djomo AN, Chimi CD. Tree allometric equations for estimation of above, below and total biomass in a tropical moist forest: Case study with application to remote sensing. *For. Ecol. Manag*. 2017; 391: 184-193. <https://doi.org/10.1016/j.foreco.2017.02.022>
23. Awé DV, Noiha NV, Zapfack L. Carbon Storage and emission factor of Savanna ecosystems in Sudano-Sahelian zone of Cameroon. *Journal of Botany Research*. 2020; 2(1): 60-67. DOI: 10.36959/771/562
24. Awé DV, Noiha NV, Zapfack L, Ali AD, Madou C. Carbon stocks in dead wood biomass of Savannah ecosystems in Northern Region Cameroon. *Journal of Botany Research*. 2019; 2(1): 60-70. DOI:10.36959/771/562
25. Awé DV, Noiha NV, Zapfack L, Vroh BTA, Nyeck B. Vegetation structure, root biomass distribution and soil carbon stock of Savannah agroecosystems in Sudano-Sahelian Zone of Cameroon. *Journal of Botany Research*. 2019; 2(1): 71-80. DOI: 10.36959/771/563
26. Awé DV, Noiha NV, Madou C, Zapfack L. Floristic composition, diversity and structure of *Khaya senegalensis* stands in Benue Department, Cameroon. *J. Trop. Resour. Sustain.Sci*. 2020; 8: 60-68. DOI:10.47253/jtrss.v8i1.165
27. Tchindebe A, Ibrahima A, Tchobsala, Mohamadou LMA. Allometric Equations for Predicting Biomass of *Daniellia oliveri* (Rolle) Hutch. & Dalz. Stands in the Sudano-Guinea Savannahs of Ngaoundere, Cameroon. *Ecology and Evolutionary Biology*. 2019; 4(2): 15-22. DOI:10.11648/j.eeb.20190402.11
28. Tchindebe A, Tchobsala, Amadou M.L.M., Ahmadou H., Adamou I. Species-Specific Allometric Equations for Predicting Biomass of *Faidherbia albida* (Del.) A. Chev. In the Sudano-sahelian Savannahs of Far-North, Cameroon. *Journal of Agriculture and Ecology Research International*. 2020; 21(6): 33-44.
29. Nogueira EM, Nelson BW, Fearnside PM. Wood density in dense forest in central Amazonia, Brazil. *For Ecol Manag*. 2005; 208: 261-286. <https://doi.org/10.1016/j.foreco.2004.12.007>
30. ASTM. Standard Test Methods for Moisture-Density (Unit Weight) Relations of Soil-Cement Mixtures ASTM D558/D558M - 19. West Conshohocken, PA: American Standards for Testing Methods. 2019; 545.
31. Picard N, Saint-André L, Henry M. Manual for building tree volume and biomass allometric equations: from field measurement to prediction. Rome : Food and Agricultural Organization of the United Nations and Montpellier, Centre de Coopération Internationale en Recherche Agronomique pour le Développement. 2012; 215.
32. Noiha NV, Awe DV, Tabue MBR, Nyeck B, Zapfack L. The Clean Development Mechanism (CDM) implementation in Africa: Allometric equation for predicting above-ground biomass in agroecosystems based on *Anacardium occidentale* L. in Sub-Saharan Africa with a case study from Cameroon. *Acta Botanica Brasilica*. (in press).
33. Biah I, Guendehou S, Goussanou C, Kaire M, Sinsin BA. Modèles allométriques pour l'estimation de stocks de biomasse dans une plantation d'anacardiés (*Anacardium occidentale* L.) au Bénin. *Bulletin de la recherche agronomique du Bénin (BRAB)*. 2018; 84: 1025-2355.
34. Flynn JJH, Holder CD. A Guide to Useful Woods of the World. 2nd ed. Forest Products Society, Madison. 2001; 618.

35. Guendehou GHS, Lehtonen A, Moudachirou M, Mäkipää R, Sinsin B. Stem biomass and volume models of selected tropical tree species in West Africa. *South Forests*. 2012; 74(2): 77–88. Available: <http://dx.doi.org/10.2989/20702620.2012.701432>
36. Chave JR, Condit S, Lao JP, Caspersen RB, Foster SP, Hubbell. Spatial and temporal variation in biomass of a tropical forest: results from a large census plot in Panama. *Journal of Ecology*. 2003 91: 240–252. <https://doi.org/10.1046/j.1365-2745.2003.00757.x>
37. Brown S, Gilespeie AJR, Lugo AE. Biomass estimation methods for tropical forest with application to forest inventory data. *Forest Sciences*. 1997; 35(4): 881–902. <https://doi.org/10.1093/forestscience/35.4.881>
38. Dong L, Zhang L, Li F. Additive biomass equations based on different dendrometric variables for two dominant species (*Larix gmelini* Rupr. and *Betula platyphylla* Suk.) in natural forests in the eastern Daxing'an Mountains, Northeast China. *Forests*. 2018; 9: 261. <https://doi.org/10.3390/f9050261>
39. Shengwang M, Quanquan J, Qijing L, Guang Z, Huimin W, Jian Y. Aboveground Biomass Allocation and Additive Allometric Models for Natural *Larix gmelinii* in the Western Daxing'anling Mountains, Northeastern China. *Forests*. 2019; 10: 150. <https://doi.org/10.3390/f10020150>
40. Mensah S, Veldtman R, Seifert T. Allometric models for height and aboveground biomass of dominant tree species in South African Mistbelt forests. *South For J For Sci*. 2016; 79: 19–30. <http://dx.doi.org/10.2989/20702620.2016.1225187>
41. Mbow C, Verstraete MM, Sambou B, Diaw AT, Neufeldt H. Allometric models for aboveground biomass in dry savanna trees of the Sudan and Sudan Guinean ecosystems of Southern Senegal. *J For Res*. 2013; 19: 340–347. <https://doi.org/10.1007/s10310-013-0414-1>
42. Thiam S, Sambou B, Mbow C, Guisse A. Élaboration de modèles allométriques d'Acacia Sénégal L. Willd pour l'analyse du carbone ligneux en milieu sahélien : cas de la zone sylvopastorale au Sénégal. *Afrique Science*. 2014; 10(3): 304–315.
43. Diédhiou I, Diallo D, Mbengue AA, Hernandez RR, Bayala R, Diémé R, Diédhiou PM, Sène A. Allometric equations and carbon stocks in tree biomass of *Jatropha curcas* L. in Senegal's Peanut Basin. *Global Ecology and Conservation*. 2017; 9: 61–69. <https://doi.org/10.1016/j.gecco.2016.11.007>
44. Mascaro J, Litton CM, Hughes F, Uowolo A, and Schnitzer SA. Is logarithmic transformation necessary in allometry? Ten, one-hundred, one-thousand-times yes. *Biological Journal of the Linnean Society*. 2014; 111: 230–233. <https://doi.org/10.1111/bij.12177>