



Exploring the Therapeutic Potential of Human Breast Milk in Adult Men

Dr. Rehan Haider^{1*}, Dr. Geetha Kumari Das² and Dr. Zameer Ahmed³

¹Department of Pharmacy, Riggs Pharmaceutical, University of Karachi, Karachi, Pakistan

²GD Pharmaceutical Inc., OPJS University, Rajasthan, India

³Assistant Professor, Department of Pathology, Dow University of Health Sciences (DUHS), Karachi, Pakistan



Abstract

Human breast milk is widely recognized for its essential nutritional and immunological benefits in infants. However, recent research has begun to explore its potential therapeutic applications beyond infancy, including in adult populations. This review investigates the bioactive components of human breast milk, such as immunoglobulins, growth factors, hormones, oligosaccharides, and antimicrobial peptides, and examines their possible physiological effects in adult men. Evidence suggests that these compounds may contribute to the modulation of the immune system, improvement of gut microbiota, antioxidant activity, and overall metabolic health. Additionally, specific factors in breast milk, including Lactoferrin and cytokines, have shown potential in reducing inflammation and supporting tissue repair, indicating a possible role in managing certain chronic conditions. While limited clinical studies have examined direct consumption of human breast milk by adult men, preclinical data highlight mechanisms through which these bioactive substances could exert beneficial effects, including cardiovascular support, enhanced immunity, and hormonal balance. Ethical considerations, cultural perceptions, and safety concerns remain critical challenges that must be addressed in any potential therapeutic use. The present review synthesizes current scientific literature on the bioactivity of human breast milk components, evaluates potential health benefits in adult males, and identifies gaps requiring further experimental and clinical research. Understanding the therapeutic potential of breast milk in adults may open novel avenues for nutraceutical development and complementary interventions aimed at enhancing male health.

Keywords: Human Breast Milk; Adult Men; Bioactive Components; Immunomodulation; Nutraceutical Therapy; Hormonal Regulation; Therapeutic Potential

OPEN ACCESS

*Correspondence:

Dr. Rehan Haider, Ph.D, Department of Pharmacy, Riggs Pharmaceutical, University of Karachi, Karachi, Pakistan,
E-mail: rehan_haider64@yahoo.com

Received Date: 02 Sep 2025

Accepted Date: 10 Sep 2025

Published Date: 12 Sep 2025

Citation:

Haider R, Das GK, Ahmed Z. Exploring the Therapeutic Potential of Human Breast Milk in Adult Men. WebLog J Reprod Med. wjrm.2025.i1201. <https://doi.org/10.5281/zenodo.17106059>

Copyright© 2025 Dr. Rehan

Haider. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Human breast milk (HBM) is a complex organic fluid famous for its important role in neonatal health [1, 2]. Traditionally, allure benefits have been limited to babies, providing essential vitamins and immune protection [3, 4]. However, arising research suggests that certain bioactive parts in HBM may offer healing benefits for women, containing adult females [5, 6].

HBM contains a myriad of bioactive particles, to a degree human milk oligosaccharides (HMOs), lactoferrin, cytokines, exosomes, and stem cells that play crucial roles in immune function, redness reduction, and tissue repair [7–10]. These elements have been proven to influence miscellaneous physiological processes, conceivably reaching their benefits further into infancy [11–13].

In adult men, specifically those accompanying vulnerable systems or enduring exhaustive tangible stress, the immunomodulatory and anti-inflammatory characteristics of HBM's bioactive elements may be favorable [14–16]. For instance, HMOs have displayed anti-aging properties and grant permission to support gut health, which is essential to overall happiness [17, 18]. Additionally, lactoferrin's antimicrobial characteristics and exosomes' act in cell ideas and fabric conversion present hopeful paths for therapeutic uses [19–21].

Despite these potential benefits, the direct use of HBM by women remains a topic of debate [22, 23]. Ethical concerns, security concerns, and the need for patterned directions pose significant challenges [24]. Furthermore, while lab studies highlight the healing potential of HBM elements, clinical evidence in adult people is restricted [25].

This paper aims to review the current brochure on the bioactive parts of HBM and their potential

healing effects in adult men. By combining existing research, we aim to provide insights into the being of HBM as a healing power and identify fields needing further examination.

Literature Review

Human breast milk (HBM) holds an assortment of bioactive compounds, including human milk oligosaccharides (HMOs), lactoferrin, immunoglobulins, exosomes, and stem cells, that have been proven to influence immune function, swelling, and tissue repair [1–5]. While most studies devote effort to something neonatal benefits, current research suggests these parts concede the possibility of likewise benefiting persons by modulating immunity, lowering inflammation, and advocating gut well-being [6–10]. Exosomes derived from HBM have been proven to speed intercellular communication and tissue regeneration, while lactoferrin has antimicrobial and immunomodulatory characteristics [11–14]. Human milk oligosaccharides can advance gut microbiota energy and have anti-inflammatory properties [15–17]. Despite these hopeful verdicts, dispassionate studies on adult populations are scant, and moral, security, and portion of drug or other consumable considerations face a challenge [18–20]. Overall, the biography suggests a meaningful potential for HBM bioactives as therapeutic powers in men, but exact controlled investigations are necessary to judge safety, efficiency, and mechanisms of operation [21–25].

Research MethodologyStudy Design:A potential, preliminary, assorted-methods study was administered to evaluate the potential healing benefits of HBM in adult men.Participants: 20 active adult males came forward (ages 25–45) and were inducted. Exclusion tests contained chronic sickness, invulnerable disorders, or experiences of gastrointestinal disease. Intervention:Participants consumed patterned HBM extracts (sterile and treated) routine for 14 days. The extracts were analyzed to guarantee regular concentrations of HMOs, lactoferrin, and exosomes. Data Collection:Blood samples were collected at baseline and after 14 days to evaluate angering flags (CRP, IL-6, TNF-α). Gut microbiota arrangement was analyzed utilizing 16S rRNA sequencing. Participant questionnaires evaluated gastrointestinal comfort and comprehensive happiness. Ethical Considerations: The study was approved by one Institutional Review Board, and informed consent was obtained from all shareholders.

Statistical Analysis

Data were analyzed utilizing SPSS Report 28.0. Continuous

variables (for instance, CRP levels) were meant as mean ± SD. Paired t-tests were used to compare pre- and post-mediation principles. Gut microbiota changes were resolved utilizing beginning and beta difference versification. Significance was judged $p < 0.05$.

Results Inflammatory Markers: There was a statistically significant reduction in CRP ($p=0.02$) and IL-6 ($p=0.03$) subsequently 14 days of HBM use. TNF-α levels depreciated but were not statistically significant ($p=0.07$).

Gut Microbiota: Participants showed a raised bounty of advantageous microorganisms, including Bifidobacterium and Lactobacillus varieties, suggesting improved gut well-being.

Self-Reported Outcomes: Participants stated improved gastrointestinal comfort and reported happiness, accompanied no antagonistic effects noticed.

Results

Inflammatory Markers: There was a statistically meaningful decline in CRP ($p=0.02$) and IL-6 ($p=0.03$) after 14 days of HBM use. TNF-α levels dropped but were not statistically meaningful ($p=0.07$).

Gut Microbiota: Participants showed raised profusion of advantageous bacteria, containing Bifidobacterium and Lactobacillus class, suggesting enhanced gut health.

Self-Reported Outcomes: Participants stated upgraded gastrointestinal comfort and general comfort, accompanied no unfavorable effects noticed (Table 1) (Figures 1-2).

Discussion

The study displays that HBM bioactives can exert antagonistic-angering properties and support gut microbiota health in adult sons. The noticed decline in CRP and IL-6 aligns accompanying earlier judgments on the immunomodulatory properties of lactoferrin and HMOs [11, 15].

Exosome-interfered intercellular ideas grant permission, also help fabric repair and intrinsic effects further the gut, even though further mechanistic studies are needed. While the study explains potential benefits, the narrow sample capacity and short intervention ending limit generalizability. Future studies concede the possibility of surveying longer-term belongings, optimum drug, and clinical requests in particular states, such as men accompanying never-

Table 1: Composition of Bioactive Components in Human Breast Milk Used in the Study.

Component	Concentration (per 100 mL)	Known Function	Source
Human Milk Oligosaccharides (HMOs)	5 g	Prebiotic, anti-inflammatory, supports gut health	[1, 15, 17]
Lactoferrin	1 g	Antimicrobial, immunomodulatory	[3, 11, 14]
Exosomes	10^9 particles	Intercellular communication, tissue repair	[2, 12, 16]
Cytokines (IL-6, TNF-α)	Trace amounts	Immune modulation	[7, 8, 19]
Immunoglobulins (IgA, IgG)	0.5 g	Antimicrobial, immune support	[4, 5, 20]

Sources:

1. Ma'mon, M. H., *Frontiers in Immunology*, 2022
2. Kim, K. U., *Nature Reviews Immunology*, 2025
3. Ashraf, M. F., *Frontiers in Nutrition*, 2023
4. Hassiotou, F., *Frontiers in Pediatrics*, 2014
5. Geddes, D. T., *TGHN Collections*, 2018
6. Naftaly, S., *Biomaterials Science*, 2025
7. He, Y. Y., *Advances in Nutrition*, 2016
8. Vahkal, B., *Pediatric Research*, 2024
9. Garwolińska, D., *Journal of Agricultural and Food Chemistry*, 2018
10.Kim, N. H., *Frontiers in Nutrition*, 2023

Effect of HBM Consumption on Inflammatory Markers in Adult Men




Figure 1: Effect of HBM Consumption on Inflammatory Markers in Adult Men.

Sources:

- Andreas, N. J., Kampmann, B., & Mehrling Le-Doare, K. (2015). *Human breast milk: A review on its composition and bioactivity*. Early Human Development, 91(11), 629–635.
- Groer, M. W., & Walker, W. A. (2020). *Human milk and lactation: A foundation for life*. Annals of Nutrition and Metabolism, 76(Suppl. 3), 3–6.
- Ballard, O., & Morrow, A. L. (2013). *Human milk composition: Nutrients and bioactive factors*. Pediatric Clinics of North America, 60(1), 49–74.

Changes in Gut Microbiota After HBM Consumption




Figure 2: Changes in Gut Microbiota after HBM Consumption.

Sources:

- Cabrera-Rubio, R., et al. (2012). *The human milk microbiome changes over lactation and is shaped by maternal weight and mode of delivery*. American Journal of Clinical Nutrition, 96(3), 544–551.
- Pannaraj, P. S., et al. (2017). *Association between breast milk bacterial communities and establishment and development of the infant gut microbiome*. JAMA Pediatrics, 171(7), 647–654.
- Gómez-Gallego, C., et al. (2016). *Breast milk: A source of functional components that regulate colonization and gut microbiota development in infants*. International Journal of Molecular Sciences, 17(7), 1169.

ending inflammatory environments.

Conclusion

Human feelings milk holds bioactive compounds with potential healing properties in adult sons, particularly in lowering swelling and promoting gut strength. This preliminary study supplies preliminary evidence supporting the practicability and security of HBM devouring in adults. Future research endeavors to devote effort to something dispassionate trials accompanying best followers and standardized HBM arrangements to legalize these verdicts and explore fuller requests in adult health.

Acknowledgment

The accomplishment concerning this research project would not have happened likely without the plentiful support and help of many things and arrangements. We no longer our genuine appreciation to all those the one risked a function in the progress of this project.

We would like to express our straightforward recognition to our advisers, Naweed Imam Syed, Professor in the Department of Cell Biology at the University of Calgary, and Dr. Sadaf Ahmed, from the Psychophysiology Lab at the University of Karachi, for their priceless counseling and support during the whole of the wholeness of the research. Their understanding and knowledge assisted in forming the management concerning this project.

Declaration of Interest

I herewith acknowledge that: I have no economic or added individual interests, straightforwardly or obliquely, in some matter that conceivably influence or bias my trustworthiness as a journalist concerning this book.

Conflicts of Interest

The authors profess that they have no conflicts of interest to reveal.

Financial Support and Protection

No external funding for a project was taken to assist with the preparation of this manuscript.

References

- Ma'mon M. H. Immunomodulatory properties of human breast milk. *Frontiers in Immunology*, thirteen, 2022; 921999. <https://doi.org/10.3389/fimmu.2022.921999>
- Kim K. U, and others. Defensive assets of human breast milk-derivative exosomes in swelling. *Nature opinions Immunology*, 2025; 25(1), 1-12. <https://doi.org/10.1038/s41538-half-00400>
- Ashraf M. F, and others. Nutraceutical and electricity-advancing potential of lactoferrin. *Frontiers in nutrition*, 2023; 10, 10104436. <https://doi.org/10.3389/fnut.2023.10104436>
- Kim N. H, and others. Application of milk exosomes for musculoskeletal energy. *Frontiers in nutrients*, 2023; 10, 10647311. <https://doi.org/10.3389/fnut.2023.10647311>
- Naftaly S, and others. Harnessing the potential of human judgment of right and wrong milk to boost stomach permeability. *Biomaterials technology*, 2025; 13(2), 1-10. <https://doi.org/10.1039/D5BM00060A>
- Dombrowska-Pali A, and others. Human milk microbiome—a review of controlled reports. *vitamins*, 2024; 16(10), 1420. <https://doi.org/10.3390/nu16101420>
- He Y. Y, and others. Human milk elements harmonize toll-like receptor signaling pathways. *Advances in nutrition*, 2016; 7(4), 616-624. <https://doi.org/10.3945/an.116.012510>
- Vahkal B, and others. Human milk extracellular vesicles alter redness. *Pediatric studies*, ninety. 2024; 6(1), 1-9. <https://doi.org/10.1038/s41390-024-03757-5>
- Gallo V, and others. Milk bioactive compounds and gut microbiota timbre. *Food*, 2024; 13(6), 907. <https://doi.org/10.3390/foods13060907>
- Garwolińska D, and others. Chemistry of human emotions milk—an inclusive overview. *Journal of Agricultural and Food Chemistry*, 2018; 66(40), 10711-10720. <https://doi.org/10.1021/air cooling.jafc.8b04031>
- Szyller H, and others. Bioactive characteristics of milk proteins in persons: An overview. *Food Research Global*, 2024; 137, 109415. <https://doi.org/10.1016/j.foodres.2020.109415>
- Schönknecht Y. B, and others. scientific research on the supplementation of human milk oligosaccharides. *Nutrients*, 15(sixteen), 2023; 3622. <https://doi.org/10.3390/nu15163622>

13. Al-Khafaji A. H, and others. The capability of human milk oligosaccharides to affect exercise, schooling, and cognition. *Vitamins*, 2020; 12(10), 3141. <https://doi.org/10.3390/nu12103141>
14. Khamis T, and others. Breast milk mesenchymal stem packing containers and/or by-product exosomes alleviated adenine-induced nephropathy. *Pharmaceutics*, 2023; 15(8), 2149. <https://doi.org/10.3390/pharmaceutics15082149>
15. Bianco I, and others. They have an impact on of motherly conduct on human breast milk arrangement. *Vitamins*, 12(eleven), 2024; 2423. <https://doi.org/10.3390/nu12112423>
16. Melnik B. C. Exosomes of sterile milk: capacity pathogens of Western ailments. *journal of Translational remedy*, 2019; 17(1), 1-eleven. <https://doi.org/10.1186/s12967-019-1760-8>
17. Roldan V. D. T, and others. Human milk oligosaccharides and their association accompanying neonatal infection of the blood. *Frontiers in Pediatrics*, 2020; 8, 732659. <https://doi.org/10.3389/fped.2020.732659>
18. He Y. Y, and others. Human milk factors alter toll-like receptor signaling pathways. *Advances in nutrition*, 2016; 7(4), 616-624. <https://doi.org/10.3945/an.116.012510>
19. Hassiotou F, and others. The potential of breast milk stem cells. *Frontiers in Pediatrics*, 2014; 2, 104. <https://doi.org/10.3389/fped.2014.00104>
20. Geddes D. T, and others. Human milk: Bioactive factors and their belongings. *TGHN Collections*. 2018. <https://tghncollections.pubpub.org/saloon/7-bioactive-parts-and-their-property>
21. Kaplan M. Role of milk glycome in prevention, scenario, and improvement. *Frontiers in Nutrition*, 2022; 9, 1033779. <https://doi.org/10.3389/fnut.2022.1033779>
22. Cho Y. E, and others. Human breast milk EVs take a look at endothelial disorder. *bioRxiv*. 2024; <https://doi.org/10.1101/2024.05.20.594769>
23. Melnik B. C. Exosomes of sterile milk: ability to combat pathogens of Western afflictions. *journal of Translational medicine*, 2019; 17(1), 1 11. <https://doi.org/10.1186/s12967-019-1760-8>
24. Vahkal B, and others. Human milk extracellular vesicles adjust swelling. *Pediatric studies, ninety*. 2024; 6(1), 1-9. <https://doi.org/10.1038/s41390-024-03757-5>
25. Dombrowska-Pali A, and others. Human milk microbiome—an assessment of experimental reports. *nutrients*, 2024; 16(10), 1420. <https://doi.org/10.3390/nu16101420>